

Beyond BMI: Muscle Mass Correlates with Reduced Low Back Pain Disability in Pre- Elderly Adults

by Susi Susanti

Submission date: 30-Jun-2025 07:16PM (UTC+0700)

Submission ID: 2708347155

File name: duced_Low_Back_Pain_Disability_in_PreElderly_Adults_copyedit.pdf (481.16K)

Word count: 3055

Character count: 17434

Beyond BMI: Muscle Mass Correlates with Reduced Low Back Pain Disability in Pre-Elderly Adults

Susi Susanti^{1*}, Resti Nurpratiwi¹, Laksmi Dewi Adzillina², Tita Rachma Ayuningtyas¹, Firdausi Kahfi Maulana¹

¹ Bachelor Program of Physiotherapy, Faculty of Medicine, Universitas Negeri Surabaya, 60213, Indonesia² Second Author ²Departement of Physiotherapy, STIKES Telogorejo Semarang, 50144, Semarang, 50144, Indonesia

*Correspondence: susisusanti@unesa.ac.id

Submitted : 27 June 2025

Accepted : 30 June 2025

Published : 30 June 2025

Abstract

Background: Low back pain (LBP) ranks among the most common musculoskeletal disorders that impact functional mobility and overall quality of life, especially in older adults who work at school teachers. Although body mass index (BMI) is frequently utilized as an indicator of health risk, it does not differentiate between fat and muscle mass, which may have varying implications for LBP risk.

Objectives: The objective of this research was to examine the correlation between body composition (BMI, muscle mass, body fat & visceral fat percentage) and the likelihood of experiencing LBP in pre-elderly individuals.

Methods: This study used a cross sectional analytic design and included 72 pre-elderly participants between 45-59 years old, who were selected through purposive sampling. Body composition metrics were obtained using a bioelectrical impedance analysis device, while LBP disability was evaluated by Oswestry Disability Index (ODI) questionnaire. The data were analyzed utilizing Spearman correlation due to the non-normal distribution of the variables with the aid of IBM SPSS Statistic 26.0 software.

Results: The analysis indicated a moderate negative correlation between muscle mass and ODI scores ($r^2 = -0.488$, $p = 0.001$), implying that higher muscle mass correlates with reduced LBP related disability. Visceral fat percentage exhibited a weak yet significant negative correlation with ODI ($r^2 = -0.293$, $p = 0.012$), whereas BMI and body fat percentage did not show significant correlation with LBP.

Conclusion: Muscle mass was identified as the most critical protective factor against LBP in pre-elderly adults, while BMI and total body fat did not serve as predictors of LBP risk. These results highlight the significance of muscle preservation in strategies aimed at preventing LBP and suggest that BMI alone is inadequate for evaluating musculoskeletal health risks in aging populations.

Keywords: body composition, low back pain, teacher school.

INTRODUCTION

⁴ Low back pain (LBP) is one of the leading causes of disability and work absence globally, particularly among individuals engaged in occupations requiring prolonged standing, repetitive movements, and static postures such as elementary school teachers (Mbarga et al., 2024; Monali K C & Saravankumar J, 2024; Vidal-Conti et al., 2021). In pre-elderly adults (ages 45-59), this condition is exacerbated by age-related physiological changes, especially in musculoskeletal and metabolic systems (Iizuka et al., 2015; Pano & Bogdani, 2021). As this age group approaches the threshold of older adulthood, the urgency to identify modifiable risk factors for LBP, including body composition changes, becomes critical for both occupational health and quality of life (Babaei Bonab, 2022; Irandoust & Taheri, 2015).

¹⁵ Recent studies have emphasized the significant relationship between altered body composition and the risk of LBP. Visceral obesity and sarcopenic conditions are increasingly recognized as contributors to chronic musculoskeletal pain, including in the lower back (Chen et al., 2023; Sakai et al., 2022). For instance, that excessive visceral fat was associated with increased musculoskeletal pain intensity (Shim et al., 2023). This study observed a strong link between LBP and sarcopenia in older adults (Roongsaiwatana et al., 2023). Supporting this, study result identified that higher body fat percentage and lower muscle mass significantly influenced LBP occurrence in middle-age populations (Jeon & Kim, 2018; Kook et al., 2023).

However, existing literature has largely focused on elderly populations or generalized working adults, with limited attention to the pre-elderly demographic, particularly those in physically and cognitively demanding professions like teaching. The cumulative impact of standing for extended hours, minimal ergonomic support, and reduced physical activity outside of work places teachers at high risk of both altered body composition and LBP. As highlighted, sarcopenia obesity can be particularly detrimental when combined with occupational stressors, yet few studies have specifically examined this interaction in pre-elderly educators (Cankurtaran et al., 2021; Sakai et al., 2023).

⁹ This study aims to explore the relationship between various body composition components, such as BMI, body fat percentage, visceral fat, and muscle mass for the risk of LBP among pre-elderly elementary school teachers. By focusing on this specific population and integrating multidimensional body composition assessment, this research offers a novel approach to identifying early intervention.

METHODS

1 Study Design and Participants

This investigation utilized a cross-sectional analytical design to assess the correlation between body composition and the risk of LBP in pre-elderly adults. This design is appropriate for identifying associations between variables without requiring long-term follow up. It also efficient for exploring potential risk factors in specific population and help generate hypotheses for future longitudinal or experimental studies. The data collection was conducted in November-December 2024 at Surabaya State Elementary School. A total of 72 participants (36 Female and 36 Male) were engaged in this research endeavor. The calculation of the sample size of this study is based on research results and calculation of the correlation analysis formula. The Participants were selected using purposive sampling technique are justified based on the need for inclusion and exclusion criteria to target population at high risk for occupational LBP, with inclusion criteria being adults 45 to 59 years old, not currently diagnosed with several neurological or orthopedic disorders, and capable of independent ambulation. Exclusion criteria included individuals with recent spinal surgery, malignancies or pregnancy.

Research Instruments

We used Oswestry Disability Index instrument for a questionnaire used to assess the level of disability related to LBP, specifically in 10 activities of daily living (Cook et al., 2021; Jenks et al., 2022; Koivunen et al., 2024). Meanwhile, for body composition measurements such as BMI (body mass index), muscle mass, body fat and visceral fat percentage using Tanita TBF-400 Total Body Composition Analyzer.

1 Data Analysis

The data were analyzed using IBM SPSS Statistics 26.0. The result of the data is non-normal distribution ($p < 0.05$), the Spearman rank correlation test was used to evaluated the relationship between body composition components (BMI, body fat & visceral fat percentage, and muscle mass) and the risk of LBP.

8 RESULTS

The results of the Spearman correlation analysis are presented in table 1. The analysis revealed that muscle mass had the strong and most significant correlation with ODI scores, showing a moderate negative relationship ($r_s = -0.488$, $p = 0.001$). This indicated that individuals with greater muscle mass tend to report lower levels of disability due to LBP. This finding

supports the hypothesis that muscular strength and tissue support around the spine may play a protective role against spinal pain and functional limitation.

Table 1. Spearman Correlation Between Body Composition Variables and ODI Score

Variable	Correlation Coefficient (<i>r</i>)	<i>p</i> -Value
BMI	0.000	1.000
Body Fat (%)	0.181	0.127
Visceral Fat (%)	-0.293*	0.012
Muscle Mass	-0.488**	0.001**

Note. *significant at $p < 0.01$, **significant at $p < 0.05$

In addition, visceral fat also showed a statistically significant negative correlation with ODI ($r = -0.293$, $p = 0.012$). While the strength of this relationship is weaker than that muscle mass, it still suggests that higher visceral fat levels are associated with reduced disability. On the other hand, body fat percentage showed a weak positive correlation with ODI ($r = 0.181$), but this relationship was not statistically significant ($p = 0.127$). BMI demonstrated no correlation with ODI scores ($r = 0.000$, $p = 1.000$). This result emphasized the limitation of BMI as an overall measure of body composition, as it does not distinguish between fat mass and lean mass. It reinforces the value of analyzing specific body composition components rather than solely on BMI when studying musculoskeletal conditions.

DISCUSSION

According to the findings of this research, the noteworthy inverse relationship between skeletal muscle mass and disability due to LBP corroborates previous research indicating the greater skeletal muscle mass correlates with diminished physical impairments and enhanced spinal functionality in elderly population (Almansour et al., 2024; Tesfaye et al., 2023). Additionally, existing literature suggests that sarcopenia, or the decline in lean body mass, considerably heightens the likelihood of disability associated with LBP, especially in conjunction with obesity (Kook et al., 2023). While our analysis revealed no significant correlation between BMI and body fat, this observation is in agreement with studies that propose BMI as an inadequate measure for predicting LBP, as it does not effectively differentiate between adipose tissue and muscle mass.

This research that greater muscle mass correlates with diminished levels of LBP disability aligns with wider patterns observed in occupational and aging studies, which underscore musculoskeletal strength as a vital protective element against LBP, especially in group subjected

to repetitive strain or extended periods of static posture (Raizah et al., 2023; Tai et al., 2019). Further highlight the importance of muscle quality and conditioning in alleviating mechanical stress on the spine and in thwarting the progression of chronic pain in at risk adults (Mushtaq et al., 2022; Peranson et al., 2024).

This study has identified muscle mass as a crucial protective factor against LBP in pre-elderly adults. However, future investigations should delve into the casual mechanism that supported this relationship through longitudinal or intervention methodologies. Furthermore, subsequent studies include measures of muscle function such as strength or power, as existing evidence indicates that muscle quality may serve as a more accurate predictor of LBP than mass alone (Imagama et al., 2019; Park et al., 2023).

CONCLUSION

This research concludes that muscle mass is the primary body composition factor linked to a decreased risk of LBP in pre-elderly adults. In contrast, BMI and body fat percentage did not exhibit significant associations, underscoring their limited diagnostic utility for LBP. Notably, visceral fat percentage was found to have a weak and inverse relationship with LBP, suggesting the need for the future research. These result underline the critical role of maintaining muscle mass in strategies aimed at preventing and managing LBP

ACKNOWLEDGMENTS

The authors would like to express their sincere gratitude to all participants who voluntarily in this study. The authors also extend theirs thanks to the research assistants and field enumerators who contribute to data collection and technical support. Appreciation is also given to the teachers at Surabaya State Elementary School for participants and Faculty of Medicine, Universitas Negeri Surabaya for administrative.

CONFLICT OF INTEREST

The author hereby declares that this research is free from conflicts of interest with any party.

AUTHOR'S CONTRIBUTION

Susanti contributes to the study concept and design, data analysis, interpretation of result and manuscript writing as the corresponding author. Nurpratiwi assisted in data collection,

literature review and statistical analysis. Adzillina contributed to the development of research instruments and supported participant recruitment and data management. Ayuningtyas was involved in formatting, writing and critical manuscript revision. Maulana contributed to data entry, visualization of findings and final manuscript proofreading.

FUNDING/SPONSORSHIP

This research does not receive external funding.

References

- Almansour, A. H., Almutairi, D. S., Alaskar, T. M., Kalkatawi, M. S., Aljubair, M. F., Alotaibi, R. S., AlHarkan, K. S., Alqahtani, H. A., & AlOmar, R. S. (2024). Prevalence of low back pain and disability among secondary school teacher in the eastern province of the Kingdom of Saudi Arabia: a cross-sectional analytical study. *Frontiers in Public Health, 12*. <https://doi.org/10.3389/fpubh.2024.1307845>
- Babaci Bonab, S. (2022). The effect of eight weeks of aqua therapy on back pain, body composition and motor function of inactive elderly women. *Scientific Journal of Rehabilitation Medicine, 10*(4). <https://doi.org/10.32598/sjrm.10.4.17>
- Cankurtaran, D., Yigman, Z. A., & Umay, E. (2021). Factors associated with paravertebral muscle cross-sectional area in patients with chronic low back pain. *Korean Journal of Pain, 34*(4). <https://doi.org/10.3344/KJP.2021.34.4.454>
- Chen, J., Wang, X., & Xu, Z. (2023). Sarcopenia and Chronic Pain in the Elderly: A Systematic Review and Meta-Analysis. In *Journal of Pain Research* (Vol. 16). <https://doi.org/10.2147/JPR.S435866>
- Cook, C. E., Garcia, A. N., Wright, A., Shaffrey, C., & Gottfried, O. (2021). Measurement Properties of the Oswestry Disability Index in Recipients of Lumbar Spine Surgery. *Spine, 46*(2). <https://doi.org/10.1097/BRS.0000000000003732>
- Iizuka, Y., Iizuka, H., Mieda, T., Tajika, T., Yamamoto, A., Ohsawa, T., Sasaki, T., & Takagishi, K. (2015). Association between neck and shoulder pain, back pain, low back pain and body composition parameters among the Japanese general population Epidemiology of musculoskeletal disorders. *BMC Musculoskeletal Disorders, 16*(1). <https://doi.org/10.1186/s12891-015-0759-z>
- Imagama, S., Ando, K., Kobayashi, K., Seki, T., Hamada, T., Machino, M., Ota, K., Tanaka, S., Morozumi, M., Kanbara, S., Ito, S., Ishiguro, N., & Hasegawa, Y. (2019). The Relationship between Neuropathic Pain and Spinal Alignment: Independent Risk Factors for Low Quality of Life in Middle-Aged and Elderly People. *Spine, 44*(19). <https://doi.org/10.1097/BRS.0000000000003073>
- Irandoost, K., & Taheri, M. (2015). The effects of aquatic exercise on body composition and nonspecific low back pain in elderly males. *Journal of Physical Therapy Science, 27*(2). <https://doi.org/10.1589/jpts.27.433>
- Jenks, A., Hoekstra, T., van Tulder, M., Ostelo, R. W., Rubinstein, S. M., & Chiarotto, A. (2022). Roland-Morris Disability Questionnaire, Oswestry Disability Index, and Quebec Back Pain Disability Scale: Which Has Superior Measurement Properties in Older Adults With Low Back Pain? *Journal of Orthopaedic and Sports Physical Therapy, 52*(7). <https://doi.org/10.2519/jospt.2022.10802>

- Jeon, K., & Kim, D. Il. (2018). The association between low body weight and scoliosis among Korean elementary school students. *International Journal of Environmental Research and Public Health*, *15*(12). <https://doi.org/10.3390/ijerph15122613>
- Koivunen, K., Widbom-Kolhanen, S., Perna, K., Arokoski, J., & Saltychev, M. (2024). Reliability and validity of Oswestry Disability Index among patients undergoing lumbar spinal surgery. *BMC Surgery*, *24*(1). <https://doi.org/10.1186/s12893-023-02307-w>
- Kook, M., Kim, I., Seo, J., Kim, H., Nam, H., & Han, N. (2023). The Relationship Between Low Back Pain and Sagittal Spinal Alignment and Back Muscle Mass in Korean Fishery Workers. *Annals of Rehabilitation Medicine*, *47*(6). <https://doi.org/10.5535/ARM.23075>
- Mbarga, J., Pichonnaz, C., Foley, R. A., & Ancey, C. (2024). Representations of physiotherapy and physiotherapists: A journey into the experience of people with chronic low back pain. *Kinesitherapie*, *24*(266). <https://doi.org/10.1016/j.kine.2023.07.009>
- Monali K C, & Saravankumar J. (2024). A Comparative Study on the Effects of Aerobic Exercises and Extension Exercises in Teachers with Low Back Pain. *Indian Journal of Physiotherapy & Occupational Therapy - An International Journal*, *18*. <https://doi.org/10.37506/exy5pj50>
- Mushtaq, M., Qamar, S., Rafiq, S., Zia, S. Z., Mohiuddin, M., Ghafoor, A., Saeed, M., & Iqbal, S. (2022). Prevalence of Low Back Pain and Its Intensity Among Teachers of Universities in Lahore. *Pakistan Journal of Health Sciences*. <https://doi.org/10.54393/pjhs.v3i06.309>
- Pano, G., & Bogdani, A. (2021). Effects of functional and traditional training on body composition and chronic low back pain treatment: Case study. *Journal of Human Sport and Exercise*, *16*(Proc3). <https://doi.org/10.14198/jhse.2021.16.Proc3.55>
- Park, M. W., Park, S. J., & Chung, S. G. (2023). Relationships Between Skeletal Muscle Mass, Lumbar Lordosis, and Chronic Low Back Pain in the Elderly. *Neurospine*, *20*(3). <https://doi.org/10.14245/ns.2346494.247>
- Peranson, J., Weis, C. A., Slater, M., Plener, J., & Kopansky-Giles, D. (2024). An interprofessional approach to collaborative management of low-back pain in primary care: A scholarly analysis of a successful educational module for prelicensure learners. *Journal of Chiropractic Education*, *38*(1). <https://doi.org/10.7899/jce-22-24>
- Raizah, A., Alzahrani, F., Albarqi, B., Abusaq, I., Alqarni, H., Alyami, I., Ahmad, I., & Reddy, R. S. (2023). Treatment Beliefs and Practices towards Low Back Pain among Teachers in Asir Region, Saudi Arabia—A Cross-Sectional Study. *Healthcare (Switzerland)*, *11*(3). <https://doi.org/10.3390/healthcare11030383>
- Roongsaiwatana, S., Sirikul, P., Chira-Adisai, W., & Warodomwicht, D. (2023). Prevalence and Associated Factors of Sarcopenia in Pre-retirement Aged With Chronic Nonspecific Low Back Pain. *Ramathibodi Medical Journal*, *46*(3). <https://doi.org/10.33165/rmj.2023.46.3.264121>
- Sakai, Y., Wakao, N., Matsui, H., Watanabe, T., Iida, H., & Watanabe, K. (2022). Clinical characteristics of geriatric patients with non-specific chronic low back pain. *Scientific Reports*, *12*(1). <https://doi.org/10.1038/s41598-022-05352-2>
- Sakai, Y., Watanabe, T., Wakao, N., Matsui, H., Osada, N., Kaneko, R., & Watanabe, K. (2023). Skeletal Muscle and Fat Mass Reflect Chronic Pain in Older Adult. *Gerontology and Geriatric Medicine*, *9*. <https://doi.org/10.1177/23337214231190146>
- Shim, J. G., Ryu, K. H., Cho, E. A., Ahn, J. H., Park, J., Lee, H. W., Kang, S., Han, S. Y., & Lee, S. H. (2023). Association between Body Composition and Chronic Low Back Pain in Korean Adults Aged over 50 Years: The Korea National Health and Nutrition Examination Survey (KNHANES) 2010–2011. *Medical Principles and Practice*, *32*(4–5). <https://doi.org/10.1159/000533354>

- Tai, K. L., Ng, Y. G., & Lim, P. Y. (2019). Systematic review on the prevalence of illness and stress and their associated risk factors among educators in Malaysia. *PLoS ONE*, *14*(5). <https://doi.org/10.1371/journal.pone.0217430>
- Tesfaye, A. H., Abere, G., Mekonnen, T. H., Jara, A. G., & Aragaw, F. M. (2023). A systematic review and meta-analysis of low back pain and its associated factors among school teachers in Africa. *BMC Musculoskeletal Disorders*, *24*(1). <https://doi.org/10.1186/s12891-023-06633-1>
- Vidal-Conti, J., Carbonell, G., Cantalops, J., & Borràs, P. A. (2021). Knowledge of low back pain among primary school teachers. *International Journal of Environmental Research and Public Health*, *18*(21). <https://doi.org/10.3390/ijerph182111306>

Beyond BMI: Muscle Mass Correlates with Reduced Low Back Pain Disability in Pre-Elderly Adults

ORIGINALITY REPORT

14%	13%	7%	2%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	www.frontiersin.org Internet Source	2%
2	jse.rezkimedia.org Internet Source	2%
3	www.researchsquare.com Internet Source	1%
4	www.bartleby.com Internet Source	1%
5	www.mdpi.com Internet Source	1%
6	www.allbusiness.com Internet Source	1%
7	fnur.stafpu.bu.edu.eg Internet Source	1%
8	ijnhs.net Internet Source	1%
9	karger.com Internet Source	1%
10	orca.cardiff.ac.uk Internet Source	1%
11	www.spoonfashion.com Internet Source	1%
12	www.scielo.br Internet Source	<1%

13 Mhlangabezi Slayi, Zuko Mpisana. "Prevalence and Diversity of Gastrointestinal Parasites and Tick Species in Communal Feedlots Compared to Rural Free-Grazing Cattle in the Eastern Cape Province, South Africa", *Parasitologia*, 2025

Publication

<1 %

14 Nader Tarabeih, Youssef Masharawi, Adel Shalata, Orabi Higla, Alexander Kalinkovich, Gregory Livshits. "Scoliosis and skeletal muscle mass are strongly associated with low back pain-related disability in humans: An evolutionary anthropology point of view", *American Journal of Human Biology*, 2022

Publication

<1 %

15 Tom P. Walsh, John B. Arnold, Angela M. Evans, Alison Yaxley, Raechel A. Damarell, E. Michael Shanahan. "The association between body fat and musculoskeletal pain: a systematic review and meta-analysis", *BMC Musculoskeletal Disorders*, 2018

Publication

<1 %

16 [bmcmusculoskeletdisord.biomedcentral.com](https://www.bmcmusculoskeletdisord.biomedcentral.com)
Internet Source

<1 %

17 eprints.uny.ac.id
Internet Source

<1 %

18 nottingham-repository.worktribe.com
Internet Source

<1 %

19 oem.bmj.com
Internet Source

<1 %

20 pubmed.ncbi.nlm.nih.gov
Internet Source

<1 %

21 rua.ua.es
Internet Source

<1%

22

Walaa Abu-Taleb, Abeer Abdelrahman Yamany, Yasser M. Aneis, Shimaa T. Abu El Kasem. "Effect of adding global postural reeducation to kendall exercises for treating asymptomatic forward head posture: A single-blinded randomized controlled trial", Journal of Bodywork and Movement Therapies, 2025

<1%

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On